

LONG-TERM EFFECTS OF INORGANIC FERTILIZERS AND FYM ON SOIL CHEMICAL PROPERTIES AND YIELD OF WHEAT UNDER RICE-WHEAT CROPPING SYSTEM

M.K. Bhatt^{1*}, R. Labanya¹, H.C. Joshi², N. Pareek¹, R. Chandra¹ and K.P. Raverkar¹

¹Department of Soil Science, College of Agriculture, G.B.P.U.A &T Pantnagar, U.S. Nagar, Uttarakhand, India

²Department of Plant Physiology, College of Basic Sciences and Humanities, G.B.P.U.A &T Pantnagar, U.S. Nagar, Uttarakhand, India

*Correspondence: mbhatt95@gmail.com

ABSTRACT

A field experiment with wheat at the Crop Research Centre, Department of Agronomy, Govind Ballabh Pant University of Agriculture and Technology, Pantnagar, under a rice–wheat system was conducted to study the long-term effect of inorganic fertilizer and FYM on physico-chemical properties of soil after 29 cycles of wheat under rice–wheat cropping system. The experiment comprised 14 treatment combinations viz., N, P and K fertilizer application as individual component and use of chemical fertilizers in conjunction with farm yard manure. The experimental treatments were replicated four times. Combined application of inorganic fertilizer and FYM resulted in a positive influx of nutrients by increasing soil pH, electrical conductivity, cation exchange capacity, organic carbon content, available nitrogen, available phosphorus, available potassium and yield of wheat in both the surface and sub-surface layer of soil. Lowest physico-chemical properties were recorded in control in both the surface and sub-surface layer while, the highest were recorded due to combined application of inorganic fertilizer along with organic manure. The highest EC (0.33 and 0.30 dsm^{-1}) was recorded under $\text{N}_{120}+\text{P}_{40}+\text{K}_{40}+\text{FYM}$ while, the highest CEC (35.30 and 28.20 $\text{Cmol}(\text{p}+)\text{kg}^{-1}$), organic carbon (1.20 and 0.80 %), available N (282.75 and 179.56 kg ha^{-1}), and P (28.31 and 12.72 kg ha^{-1}) was recorded due to $\text{N}_{180}+\text{P}_{80}+\text{K}_{40}+\text{Zn}(\text{F})+\text{FYM}$. The highest available K (223.45 and 109.33 kg ha^{-1}) was recorded due to $\text{N}_{120}+\text{P}_{40}+\text{K}_4+\text{FYM}$. The lowest yields (grain, straw, and biological) were recorded in control while, the highest crop yields were recorded under $\text{N}_{180}+\text{P}_{80}+\text{K}_{40}+\text{Zn}(\text{F})+\text{FYM}$. The present investigation clearly points out the significance of balanced and imbalanced use of nutrients including FYM in rice-wheat cropping system for improving the various physico-chemical soil quality indices and yield of wheat crop over a long period.

Keywords: Long term fertilizer experiment, Wheat, FYM, Chemical indices, Soil quality.

INTRODUCTION

Rice and wheat are both the dominant components of the Indian food security system as they are consumed as major staple food. Presently, the rice-wheat cropping system in the Indo-Gangetic plains is showing sign of 'fatigue' due to continuous cropping of this highly nutrient and water exhaustive cereal-cereal system for the last three decades. Imbalance use of chemical fertilizers alone tends to decline soil quality and fertility over a period of years with given

inputs. Therefore, the most logical way to manage long-term fertility and productivity of soil is integrated use of inorganic and organic sources of plant nutrients. For the development of sustainable food production system maintenance and management of soil fertility is pivotal (Doran *et al.*, 1988) the logical way emerging to manage long-term fertility and productivity of soil is integrated use of organic and inorganic sources of plant nutrients to address the concern of excess and/ or depletion of nutrients (Aulakh *et al.*, 2008).

An integrated use of manure with chemical fertilizer resulted in build-up of available nutrients in soil much more effectively than that of chemical fertilizer alone (Prasad *et al.*, 1995). They also suggested that use of manure holds a great promise in rice-wheat system not only for higher productivity levels but also against emergence of multiple nutrient deficiencies and deterioration of soil health. Dhiman *et al.*, 2007 reported that organic carbon, available P and K were highest where an additional manure application was made. Soil organic matter and total nitrogen levels were increased by about 120 % over 150 years in the manured plots. Such carbon stores might have appreciated the sink for global carbon (Tillman, 1998). Availability of nutrients in soil is governed by several ongoing chemical processes such as weathering and buffering, and properties like organic matter content, CEC and potential of hydrogen (pH). Larson *et al.*, 1991 chose those chemical properties for assessing soil quality that either inhibit root growth or quantitatively affect nutrient supply or modify their ability. Reganold *et al.*, 1995 used certain chemical parameters related to nutrient availability as a measure of soil quality like CEC, total N, P and pH; and extractable P, S, Ca, Mg and K. It was suggested that total and available plant nutrients and nutrient cycling rates should also be considered in soil quality measurement (Karlen *et al.*, 1992).

Under continuous cropping, changes in soil fertility and the resultant crop productivity can be related with nutrient imbalances, which have been recognized as one of the most important factors that limits crop yield. Even the applications of recommended NPK fertilizers, devoid of organic manure have not sustained soil quality. There are many reports based on the 10-15 years old experiments on the chemical and physical properties from the long term experiments in India and abroad separately, however very scanty information is available in respect of varying nutrient management practices in the rice-wheat cropping system and their impact on quality of Mollisols.

MATERIALS AND METHODS

The present investigation was carried out in an ongoing long term experiment, after twenty nine years, located at Norman. E. Borlaug Crop Research Centre, G.B. Pant University of Agriculture and Technology, Pantnagar, U. S. Nagar, Uttarakhand which was initiated in 1984. The experimental site lies in *Tarai* plains about 30 km southward of foothill of shivalik range of Himalayas at 29° N latitude, 79° 29' E

longitude and at an altitude of 243.8 m above the mean sea level. The chemical analysis of top 15 cm soil showed that it was rich in organic matter and medium in phosphorus and potassium and neutral to slightly alkaline in reaction. The field experiment has been laid in Randomized Block Design (RBD) in four replications with fourteen treatments comprising control, N, NP, NPK, varying doses alone and with FYM. The N, P and K were provided as per treatment details through urea, single super phosphate and potassium, respectively. In one treatment N and P was provided using Di-ammonium phosphate. The FYM was applied @5 tonne ha⁻¹ only in rice crop.

Collection of soil samples and processing

Individual soil samples from each plot were collected from two depths (0-15 and 15-30 cm) after harvesting of wheat crop in the year 2013-14. These individual soil samples were pooled to get one representative composite sample each for every depth from all the plots.

The pH was determined in 1:2 soils: water ratio and after half an hour of equilibrium, the pH was determined with the help of a glass electrode on microprocessor based pH meter (Jackson 1967). Electrical conductivity of the soil was measured in 1:2 soils: water suspension at 25°C with digital microprocessor based conductivity meter (Bover *et al.*, 1965) Cation exchange capacity indicates the total quantity of negative charge per unit weight of soil or colloids. The cation exchange capacity in soil was determined by ammonium acetate method (Bache 1976). The organic carbon content in soil was determined by following the modified Walkley *et al.*, 1934 method as described by Jackson 1967. Available nitrogen in soil was determined by alkaline potassium permanganate method (Subbiah *et al.*, 1956). Available phosphorus in soil was extracted by using sodium bi-carbonate extractant (0.5 M NaHCO₃) adjusted to pH 8.5 (Olsen *et al.*, 1954). Available potassium in soil mainly refers to exchangeable + water soluble K. The exchangeable K constitutes the major portion of available potassium except in saline and saline sodic soil. Available K was determined by neutral ammonium acetate (1N NH₄OAc; pH 7) method outlined by Black 1965.

Statistical analysis

The experimental data were analyzed using the statistical program STPR of G.B. Pant University of Agriculture and

Technology, Pantnagar in a Randomized Block Design. Analysis of Variance and critical difference (CD) between treatments was calculated at 5% level of significance. Correlation coefficients were computed using SPSS version 16.

RESULTS AND DISCUSSION

Soil pH

The data in respect of the pH at surface and sub-surface depth of soil as influenced due to application of nutrients in wheat crop for twenty-nine years through fertilizer and /or FYM is depicted in Table 1. The soil pH at 0-15 cm depth varied significantly after twenty-ninth cycle of rice crop however in sub-surface layer no significant impacts of fertilizer application was observed. The lowest pH in the surface soil after wheat crop was registered under control while the highest pH to the tune of 8.19 was recorded due to $N_{120}+P_{40}+K_{40}+FYM$. There was no remarkable variation in soil pH due to various treatments of nutrient management through chemical fertilizers alone or in combinations with FYM for 29 years. An application of chemical fertilizers over a long period leads in increase in pH Tyagi 1989. However, fertilizer application in conjunction with FYM reduced the impact in comparison to control. Similar findings have been recorded earlier by Sime 2001; Sharma 2004 and Bhatt 2012. It may be attributed to buffering action of soils that does not permit to alter the soil pH.

Electrical Conductivity

Data presented in Table 1 depicts that an application of nutrients through fertilizer alone and along with FYM had non-significant influence on soil EC at surface as well as sub-surface depths after harvest of wheat crop. The lowest electrical conductivity was recorded in control (0.27 dsm^{-1} and 0.24 dsm^{-1}) in both surface and sub-surface soil while the highest electrical conductivity (0.33 dsm^{-1} and 0.30 dsm^{-1}) was recorded with the treatment $N_{120}+P_{40}+K_{40}+FYM$ in surface and sub-surface soil. Addition of NPK fertilizers increases accumulation of salt concentration in soil which contributes to increased electrical conductivity of soil. A slight increase in electrical conductivity due to continuous fertilizer use under intensive cropping was reported (Sime 2001). Similar results were obtained (Singh *et al.*, 2007; Das 2007). The enhanced electrical conductivity due to FYM was attributed to the decomposition of organic matter (Larson *et al.*, 2002; Babu *et al.*, 2007).

Cation exchange capacity

An application of fertilizer nutrients alone and in combination with FYM influenced soil CEC significantly in surface soil as well as sub-surface soil after harvest of wheat (Table 1). In both surface and sub-surface soil layer lowest cation exchange capacity (27.10 and $22.50 \text{ Cmol (p}^+) \text{ kg}^{-1}$ soil) was recorded in control. On the other hand an application of $N_{180}+P_{80}+K_{40}+Zn$ (F) + FYM over a period of twenty-nine years favored the highest cation exchange capacity (35.30 and $28.20 \text{ Cmol (p}^+) \text{ kg}^{-1}$ soil) in both the surface and sub-surface soil layer. The balanced and imbalanced use of inorganic chemical fertilizers along with FYM increased the CEC of soil in comparison to control. The increase in CEC might be due to improvement in organic matter content of soil. CEC and exchangeable Ca^{2+} content of the soil improved with the application of organic sources of nutrients either alone or in combination with fertilizers when used over a long period of time. Long-term FYM/manure addition increases CEC due to the colloidal nature of organic matter (Patiram *et al.* 1993; Sinha *et al.*, 1997). The CEC was also found to be lowest in plots receiving N only and it increased with increase in soil pH. Lime and FYM additions increased the CEC and exchangeable Ca^{2+} and Mg^{2+} status of the soil due to increase in biomass production and its incorporation in the soil. A gradual increase in CEC due to graded levels of NPK was also recorded probably due to higher content of crop residues (Prasad *et al.*, 1996; Jeegdeshwari *et al.*, 2001).

Table 1. Effect of long-term fertilizer application at varying levels on soil pH, EC and CEC after twenty-ninth cycle of wheat crop at different depths under rice-wheat cropping system

Treatments		pH		EC (dSm ⁻¹)		CEC (Cmol (p ⁺) kg ⁻¹ soil)	
		0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	Control	7.94	8.18	0.27	0.24	27.10	22.50
T ₂	N ₁₂₀	8.04	8.17	0.30	0.28	28.50	23.20
T ₃	N ₁₂₀ +P ₄₀	8.15	8.22	0.28	0.26	30.30	23.70
T ₄	P ₄₀ +K ₄₀	8.17	8.17	0.29	0.27	31.00	24.40
T ₅	N ₁₂₀ +K ₄₀	8.06	8.19	0.31	0.28	31.50	25.10
T ₆	N ₁₂₀ +P ₄₀ +K ₄₀	8.18	8.21	0.32	0.29	33.20	26.80
T ₇	N ₁₂₀ +P ₄₀ +K ₄₀ +Zn(F)	8.16	8.22	0.32	0.29	33.30	26.70
T ₈	N ₁₂₀ +P ₄₀ +K ₄₀ +FYM	8.19	8.17	0.33	0.30	34.30	27.70
T ₉	N ₁₂₀ +P ₄₀ +K ₄₀ +Zn(F)+FYM	8.09	8.19	0.32	0.29	34.40	27.50
T ₁₀	N ₁₈₀ +P ₈₀ +K ₄₀ +Zn(F)+FYM	8.01	8.20	0.29	0.27	35.30	28.20
T ₁₁	N ₁₅₀ +P ₄₀ +K ₄₀	8.15	8.23	0.30	0.28	32.90	26.80
T ₁₂	N ₁₈₀ +P ₈₀ +K ₄₀ +Zn(F)	8.14	8.19	0.31	0.29	30.80	26.70
T ₁₃	N ₁₈₀ +P ₈₀ +Zn(F)	8.15	8.22	0.28	0.25	30.20	25.40
T ₁₄	N ₁₂₀ +P ₄₀ +K ₄₀ (DAP)	8.11	8.23	0.30	0.28	28.50	25.60
S. Em±		0.05	0.05	0.01	0.01	0.99	0.84
C.D. (5%)		0.15	NS	NS	NS	2.89	2.45
C.V. (%)		1.27	1.27	9	8.04	5.43	5.64

Organic carbon

An application of inorganic fertilizer at varying levels alone or in combination with FYM had significant influence on

change in soil organic carbon content after twenty-nine years of rice-wheat cropping system Table 2 and fig. 1. The soil organic carbon content, an indicator of soil fertility, was significantly influenced due to fertilizer application. No application of fertilizer/FYM significantly decreased organic carbon (0.72 and 0.58%) content in control both the surface and sub-surface soil layer while, the highest organic carbon (1.20 and 0.80 %) content in both the surface and sub-surface layer was registered due to application of N₁₈₀+P₈₀+K₄₀+ Zn(F)+FYM. The higher level of soil organic carbon content with the long-term fertilizer application along with FYM may be due to enhanced root growth, resulting more organic residue in soil, which after decomposition might have increased the soil organic carbon content. These results are in conformity with the finding of Singh *et al.*, 1999 who reported the enhanced soil organic carbon status after ten years of continuous rice-wheat cropping under varying fertilizer and manure treatment in Mollisols at Pantnagar. They observed that the status of organic matter dropped down drastically in all the treatments except in plots receiving FYM. It was ranged from 1.78% in PK to 2.09% with NPK + FYM from its initial value of 2.10 per cent. Continuous application of fertilizers alone and in combination with graded level of FYM for soybean based cropping system was studied by Babhulkar *et al.*, 2000 and reported that highest organic C status (6.2 g kg⁻¹) under the application of higher rate of FYM (7.5 Mg ha⁻¹) with half dose of N and P fertilizer. The maximum increase in soil organic carbon content was observed with integrated use of inorganic fertilizers (N+P+K) and organic manure. The increase in soil organic carbon content with the long-term use of manure in rice-wheat (Beri *et al.*, 1995).

Available N

Available nitrogen content in soil as affected by continuous application of balanced and imbalanced fertilization with or without FYM in rice-wheat system over a period of twenty-nine years is summarized (Table 2 and Fig 2). All the fertilizer treatments resulted in significantly higher available nitrogen in soil, after harvest of rice over control. The lowest available nitrogen was observed in control (151.88 and 111.08 kg ha⁻¹) in both the surface and sub-surface soil, while the highest value of available nitrogen (282.75 and 179.56 kg ha⁻¹) in surface and sub-surface was recorded due to N₁₈₀+P₈₀+K₄₀+Zn(F)+FYM. Continuous cropping of rice-wheat system for twenty-nine years increased the available N significantly with the use of fertilizer N in combination with P and K fertilizer, as well as with combined use of inorganic

fertilizer and FYM. The increase in available nitrogen content in soil due to application of FYM may be attributed to the mineralization of nitrogen and greater multiplication of soil microbes which converts organically bound N into inorganic form. The lowest value of available N in control may be due to mining of nutrients with continuous cropping without fertilization over the years. Application of fertilizers either alone NPK or in conjunction with organics were significantly superior to control. Significantly higher available N in fallow, NPK and N alone plots over control and in 100% NPK + FYM as compared to 100% NPK in silty clay loam at Pantnagar after 6 year of intensive cropping was reported Bhardwaj 1983; Singh *et al.*, 1985). Increased available N after 4-year application of FYM and green manuring in soil may be due to early decomposition of green manure crops vis-a-vis FYM Debnath *et al.*, 1972). In a long-term fertilizer experiment under rice-wheat system significant effect of fertilizer treatment on available nitrogen content after wheat in both the soil depths was observed (Bhatt *et al.*, 2012). Available nitrogen was found lower in deeper soil layer (15-30 cm) than in surface layer (0-15 cm) in both the years. Highest available nitrogen in both the profile depth was with 100% NPK+FYM, which was significantly higher than all the applied nutrient treatments except 150% NPK (Bhatt 2012).

Table 2. Effect of long-term fertilizer application at varying levels on soil organic carbon and available nitrogen after twenty-ninth cycle of wheat crop at different depths under rice-wheat cropping system

	Treatments	Organic C (%)		Available N (kg ha ⁻¹)	
		0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	Control	0.72	0.58	151.88	111.08
T ₂	N ₁₂₀	0.81	0.63	189.54	156.75
T ₃	N ₁₂₀ +P ₄₀	0.85	0.66	162.91	139.32
T ₄	P ₄₀ +K ₄₀	0.79	0.69	173.84	146.75
T ₅	N ₁₂₀ +K ₄₀	0.83	0.70	185.68	150.46
T ₆	N ₁₂₀ +P ₄₀ +K ₄₀	0.86	0.74	195.81	160.75
T ₇	N ₁₂₀ +P ₄₀ +K ₄₀ +Zn(F)	0.89	0.76	210.88	165.88
T ₈	N ₁₂₀ +P ₄₀ +K ₄₀ +FYM	1.14	0.76	252.88	175.29
T ₉	N ₁₂₀ +P ₄₀ +K ₄₀ +Zn(F)+FYM	1.17	0.79	248.20	164.43
T ₁₀	N ₁₈₀ +P ₈₀ +K ₄₀ +Zn(F)+FYM	1.20	0.80	282.75	179.56
T ₁₁	N ₁₅₀ +P ₄₀ +K ₄₀	0.84	0.72	243.10	160.91
T ₁₂	N ₁₈₀ +P ₈₀ +K ₄₀ +Zn(F)	0.87	0.75	236.02	158.74
T ₁₃	N ₁₈₀ +P ₈₀ +Zn(F)	0.83	0.70	241.51	159.91
T ₁₄	N ₁₂₀ +P ₄₀ +K ₄₀ (DAP)	0.78	0.64	192.67	150.77
	S.Em±	0.04	0.03	6.29	4.91
	C.D. (5%)	0.10	0.08	18.05	14.10
	C.V. (%)	7.72	7.73	5.93	6.30

Available P

The data pertaining to available phosphorus in soil as influenced by continuous balanced and imbalanced fertilization in rice-wheat system are summarized in Table 3

and fig. 3. The fertilizer treatments had significant influence on available soil phosphorus after harvest of wheat crop. Highest amount of available phosphorus (28.31 and 12.72 kg ha⁻¹) in surface as well as sub-surface soil was recorded under N₁₈₀+P₈₀+K₄₀+Zn(F)+FYM while, the lowest available phosphorus (15.14 kg ha⁻¹ and 7.02 kg ha⁻¹) was observed in control both the surface and sub-surface soil. Continuous cropping with combined application of inorganic fertilizers along with FYM increased the available P in soil. In general, the combined application of organics with fertilizers resulted in higher available P content over application of inorganic fertilizers alone. Build-up in available P with the conjoint use of fertilizers and organics could be ascribed to the release of organic acids during decomposition which in turn helped in releasing native phosphorus through solubilizing action of these acids. Also, organic matter forms a coating on sesquioxides and makes them inactive and thus reduces the phosphate fixing capacity of soil, which ultimately helps in release of ample quantity of plant available P. The similar results were obtained by (Gangwar *et al.*, 1996; Bhatt 2012; Sharma *et al.*, 2014).




Fig.1. Effect of long-term fertilizer application at varying levels on soil organic carbon after twenty-ninth cycle of wheat crop at different depths under rice-wheat cropping system




Fig. 2. Effect of long-term fertilizer application at varying levels on available nitrogen after twenty-ninth cycle of wheat crop at different depths under rice-wheat cropping system

Available K

Overall treatment effect on available potassium was statistically significant in surface and sub-surface soil after harvest of rice crop Table 3 and fig 4. The treatments which do not received potassium fertilizer *viz.* N₁₂₀+P₄₀, N₁₂₀ and control had significantly lower amount of available soil potassium than the potassium consisting treatments. The lowest available potassium (168.13 kg ha⁻¹ and 82.85 kg ha⁻¹) in soil was observed in N₁₂₀ in both the surface and sub-surface soil layer while the highest amount of available potassium in surface and sub-surface soil layer (223.45 kg ha⁻¹ and 109.33 kg ha⁻¹) was measured under (N₁₂₀+P₄₀+K₄₀+FYM). Increase in available K in T₈, T₉ and T₁₀ treatments due to combined use of inorganic fertilizers along with FYM may be ascribed to the reduction of K fixation and release of K due to interaction of organic matter with clays, besides the direct K addition to the soil (Urkurkar *et al.*, 2010; Subehia *et al.*, 2012). Shri Ram *et al.*, 2016 found that continuous application of inorganic fertilizer, FYM and cropping for 41 years significantly increased available P content and the greatest available K was recorded under the FYM treated plots followed by super-optimal dose of NPK fertilizers. Respectively, the lowest value of available K in control may be due to mining of nutrients with continuous cropping without fertilization over the years. The treatment wise trends in sub-surface layer were almost similar as was observed under surface layer. However, the availability of potassium was decreased with increase in depth. The higher amount of K in the surface layer could be attributed to more exposure of K bearing minerals to weathering and or upward translocation of K from sub-surface layer by capillary rise or due to addition of K through plant residue, manure and fertilizers (Rao *et al.*, 2013).

Table 3. Effect of long-term fertilizer application at varying levels on soil available phosphorus and available potassium after twenty-ninth cycle of wheat crop at different depths under rice-wheat cropping system

Treatments		Available P (kg ha ⁻¹)		Available K (kg ha ⁻¹)	
		0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	Control	15.14	7.02	180.69	88.05
T ₂	N ₁₂₀	17.07	7.19	168.13	82.85
T ₃	N ₁₂₀ +P ₄₀	19.71	9.19	168.69	83.21
T ₄	P ₄₀ +K ₄₀	22.53	10.57	205.93	95.05
T ₅	N ₁₂₀ +K ₄₀	16.12	7.17	211.53	102.8
T ₆	N ₁₂₀ +P ₄₀ +K ₄₀	19.18	8.96	214.89	102.08
T ₇	N ₁₂₀ +P ₄₀ +K ₄₀ +Zn(F)	18.50	8.09	205.09	99.96
T ₈	N ₁₂₀ +P ₄₀ +K ₄₀ +FYM	27.64	12.59	223.45	109.33
T ₉	N ₁₂₀ +P ₄₀ +K ₄₀ +Zn(F)+FYM	27.10	12.05	221.33	102.67
T ₁₀	N ₁₈₀ +P ₈₀ +K ₄₀ +Zn(F)+FYM	28.31	12.72	212.65	101.49
T ₁₁	N ₁₅₀ +P ₄₀ +K ₄₀	18.77	8.65	214.05	102.01
T ₁₂	N ₁₈₀ +P ₈₀ +K ₄₀ +Zn(F)	25.62	11.38	205.93	98.69
T ₁₃	N ₁₈₀ +P ₈₀ +Zn(F)	20.16	9.42	203.69	93.09
T ₁₄	N ₁₂₀ +P ₄₀ +K ₄₀ (DAP)	17.02	8.02	182.41	90.29
S.Em±		0.90	0.35	5.80	4.09
C.D. (5%)		2.59	0.99	16.66	11.74
C.V. (%)		8.62	7.28	5.76	8.47




Fig. 3. Effect of long-term fertilizer application at varying levels on soil available phosphorus after twenty-ninth cycle of wheat crop at different depths under rice-wheat cropping system




Fig. 4. Effect of long-term fertilizer application at varying levels on available potassium after twenty-ninth cycle of wheat crop at different depths under rice-wheat cropping system

Yields

The fertilizer treatments comprising fertilizer alone or with FYM had significant effect on grain yield of wheat Table 4 and fig 5. The results obtained indicated that the highest grain yield (3524 kg ha^{-1}) was recorded due to application of $\text{N}_{180}+\text{P}_{80}+\text{K}_{40}+\text{Zn}(\text{F}) + \text{FYM}$ while, the lowest grain yield (859 kg ha^{-1}) was recorded under control. The results obtained indicated that the highest straw yield was recorded due to application of $\text{N}_{180}+\text{P}_{80}+\text{K}_{40}+\text{Zn}(\text{F}) + \text{FYM}$ while, the lowest straw yield was recorded under control. The highest biological yield was recorded with $\text{N}_{180}+\text{P}_{80}+\text{K}_{40}+\text{Zn}(\text{F}) + \text{FYM}$ while, the lowest biological yield was observed under control. The fertilizer treatments had significant influence on grain, straw and biological yields. The addition of FYM along with inorganic fertilizers produced higher grain yield of wheat. It could be due to additional supply and availability of nutrients through FYM on the one hand and improved physical, chemical and biological condition of soil on the

other hand. These findings are in conformity with the findings of Sharma *et al.*, 1999; Singh *et al.*, 1999; Yaduvanshi 2003. Combination of NPK along with FYM and Zn increased the grain yield of both the crops because of attaining better fertility status resulting in good supply of nutrients. Such inference has also been drawn from several long-term fertility experiments conducted all over India (Nambiar *et al.*, 1989; Dawe *et al.*, 2000; Yadav 2009).

Table 4. Effect of long-term fertilizer application at varying levels on yield of rice and wheat after twenty-ninth cycle under rice-wheat cropping system

Treatments		Wheat (kg ha^{-1})		
		Mean grain yield	Mean straw yield	Biological yield
T ₁	Control	859	1196	2056
T ₂	N_{120}	1114	1531	2645
T ₃	$\text{N}_{120}+\text{P}_{40}$	2109	2937	5046
T ₄	$\text{P}_{40}+\text{K}_{40}$	1432	2052	3485
T ₅	$\text{N}_{120}+\text{K}_{40}$	1259	1745	3004
T ₆	$\text{N}_{120}+\text{P}_{40}+\text{K}_{40}$	2534	3331	5866
T ₇	$\text{N}_{120}+\text{P}_{40}+\text{K}_{40}+\text{Zn}(\text{F})$	2951	3928	6880
T ₈	$\text{N}_{120}+\text{P}_{40}+\text{K}_{40}+\text{FYM}$	3264	4362	7626
T ₉	$\text{N}_{120}+\text{P}_{40}+\text{K}_{40}+\text{Zn}(\text{F})+\text{FYM}$	3429	4526	7956
T ₁₀	$\text{N}_{180}+\text{P}_{80}+\text{K}_{40}+\text{Zn}(\text{F})+\text{FYM}$	3524	4827	8352
T ₁₁	$\text{N}_{150}+\text{P}_{40}+\text{K}_{40}$	2596	3555	6150
T ₁₂	$\text{N}_{180}+\text{P}_{80}+\text{K}_{40}+\text{Zn}(\text{F})$	2830	3907	6738
T ₁₃	$\text{N}_{180}+\text{P}_{80}+\text{Zn}(\text{F})$	3090	4285	7375
T ₁₄	$\text{N}_{120}+\text{P}_{40}+\text{K}_{40}$ (DAP)	2813	3899	6712
S.E.m±		311.8	136.5	256.9
C.D. (5%)		978.4	391.8	737.5
C.V. (%)		10.9	8.3	9.0




Fig. 5. Effect of long-term fertilizer application at varying levels on yield of wheat crop after twenty-ninth cycle under rice-wheat cropping system

CONCLUSION

Long-term use of inorganic fertilizers and organic manure (FYM) found superior in comparison to alone application of inorganic fertilizers to sustain the crop productivity and soil fertility and to enhance the soil quality in a rice-wheat cropping system grown on a Mollisol in northern India. The inorganic fertilizers along with organic manure was found to be a viable option for restoring soil organic carbon and nutrient turnover, thereby improving the availability of

nutrients in soil, maintaining soil quality, and helping achieve sustainable productivity of wheat crop for the long run under irrigated moisture regimes. Continuous application of NPK with Zn and FYM improve the yield sustainability and soil quality. Cumulative effect of long-term use of fertilizers lowered the risk of soil quality over a single year of application of fertilizers. Therefore, judicious application of inorganic and organic nutrients in an integrated manner is essential for proper nutrient supply, sustaining crop productivity and soil quality in a long-term rice-wheat cropping system.

REFERENCES

- Aulakh M, Grant CA (2008). Integrated nutrient management for sustainable crop production. *CRC Press*, 2(1): 25-39.
- Babhubhakar PS, Wandile RM, Badole WP, Balpande SS (2000). Residual effect of long-term application of FYM and fertilizers on soil properties (Vertisol) and yield of soybean. *Journal of Indian Society of Soil Science*, 48: 89-92.
- Babu M, Reddy CM, Subramanyam A, Balaguravaiah D (2007). Effect of integrated use of organic and inorganic fertilizers on soil properties and yield of sugarcane. *Journal of the Indian Society of soil Science*, 55(2):161-166.
- Bache BW (1976). The measurement of cation exchange capacity of soils. *Journal of the Science of Food and Agriculture*, 27(3): 273-80.
- Beri V, Sidhu BS, Bahl GS, Bhat AK (1995). Nitrogen and phosphorus transformations as affected by crop residue management practices and their influence on crop yield. *Soil Use and Management*, 11(2):51-54.
- Bhardwaj V (1983). Some soil properties, crop yields and nutrient uptake as affected by intensive cropping and fertilization in the long term fertilizer experiment at Pantnagar. Ph.D. Thesis. Submitted to G.B.P.U.A&T., Pantnagar, U.S. Nagar (U.K).
- Bhatt B (2012). Effect of long term fertilizer application in rice-wheat system on crop productivity and soil. Ph. D. Thesis submitted to G.B.P.U.A. & T., Pantnagar, India, 135-145.
- Black CA (1965). Methods of Soil Analysis Part 1 and 2. American Society of Agronomy, Inc.; USA, 1-12.
- Bover CA, Wilcox LA (1965). Soluble salts. In : *Method of soil Analysis*. Part-2, Chemical and Microbiological properties. *Agronomy Monograph*. ASA and SSSA, Madison, Wisconsin, USA 433-451.
- Das T (2007). Studies on long-term impact of NPK fertilizers and manure on yield, grain quality and nutrient uptake by wheat and relationship with physical and chemical properties of Mollisol. Thesis, Master of Science in agriculture (Soil Science), G.B.P.U.A. &T. Pantnagar, 12-23.
- Dawe D, Dobermann A, Moya P, Abdurachman S, Singh B, Lal P, Li SY, Lin B, Panaullah G, Sariam O, Singh Y (2000). How widespread are yield declines in long-term rice experiments in Asia. *Field Crops Research*, 66(2):175-93.
- Debnath NC, Hajra JN (1972). Transformation of organic matter and soil relation to mineralization of organic carbon and nutrient availability. *Journal of Indian Society of Soil Science*, 20: 45-102.
- Dhiman, SD, Om Hari and Singh, D (2007). Effect of long term nutrient management on soil properties and productivity in rice (*Oryza sativa*)-wheat (*Triticum aestivum*) sequence. *Ind. J. Agron*, 32: 112-120.
- Doran JW, Fraser DG, Culik MN, Liebhardt WC (1988). Influence of alternative and conventional agricultural management on soil microbial processes and nitrogen availability. *American Journal of Alternative Agriculture*, 2(3): 99-106.
- Gangwar SP, Ram N (1996). Long-term impact of fertilizers on the relationship of inorganic forms of phosphorus with its availability and uptake by rice in a hapludoll soil. *Acta Agronomica Hungarica*, 44:29-34.
- Jackson ML (1967). Prentice Hall of India. Pvt. Ltd., New Delhi. 498-506.
- Jeegadeshwari PV, Senthil Kumar PS, Kumaraswamy K (2001). Sustenance of soil fertility status of an Alfisol under fifteen years of continuous fertilization and cropping of rice. *Crop Research*, 22: 25-28.
- Karlen DL, Eash NS, Unger PW (1992). Soil and crop management effects on soil quality indicators. *American Journal of Alternative Agriculture*, 7(1-2): 48-55.
- Larson WE, Pierce FJ (1991). Conservation and enhancement of soil quality. In Evaluation for sustainable land management in the developing world: *proceedings of the International Workshop on Evaluation for Sustainable Land Management in the Developing World*, 6-24.
- Larson WE, Pierce, FJ (2002). Conservation and enhancement of soil quality. In: Evaluation of sustainable management in the developing world. *Int. Board Soil Research and Management*, Bangkok, Thailand 175-203

- Nambiar KM, Abrol IP (1989). Long term fertilizers experiments in India. LTFE Res Bull, *Indian Agril Res Bull*, New Delhi.
- Olsen SR (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. United States Department of Agriculture; Washington.
- Patiram, Singh KA (1993). Effect of continuous application of manure and nitrogenous fertilizer on some properties of acid Inceptisol. *Journal of Indian Society of Soil Science*, 41: 430-433.
- Prasad B, Prasad J, Singh K (1996). Effect of nutrient management for sustainable rice and wheat production in calcareous soil amended with green manure and Zn. *Fertilizer News* 40(2): 34-45.
- Prasad B, Sinha SK (1995). Nutrient recycling through crop residues management for sustainable rice and wheat production in calcareous soil. *Fertilizer News*, 40(1):15-28.
- Rao S, Pathak H, Ghosh AB (2013). Effect of intensive cropping and fertilizer use on the crop removal of sulphur and zinc and their availability in soil. *Fertilizer Resource*, 2: 303-08.
- Reganold JP, Palmer AS (1995). Significance of gravimetric versus volumetric measurements of soil quality under biodynamic, conventional, and continuous grass management. *Journal of Soil and Water Conservation*, 50(3): 298-305.
- Sharma RK, Chauhan DS, Nagarajan S (1999). Current Status of Crop Response to Fertilisers in Different Agro-climatic Zones-Experiences of All India Co-ordinated Wheat Improvement Project. *Fertilizer News*, 44(7): 22-34.
- Sharma U, Subehia SK (2014). Effect of long-term integrated nutrient management on rice (*Oryza sativa* L.)-wheat (*Triticum aestivum* L.) productivity and soil properties in North-Western Himalaya. *Journal of the Indian Society of Soil Science*, 62(3): 248-54.
- Sharma M (2004). Long term effect of fertilizers and manure on some physical properties of soil and nutrient uptake by wheat. Thesis Master of Science in Agriculture (Soil Science), G. B. P. U. A. & T Pantnagar, 15-20
- Sime T (2001). Long term effect of fertilizers and manures on the availability and uptake of calcium, magnesium and sulphur by wheat in a Mollisol. Thesis, Master of Science in Agriculture (Soil Science), G.B.P.U.A. & T. Pantnagar 5-19.
- Singh B, Brar SP (1985). Effect of organic manures and nitrogen on grain yield and soil properties in a maize-wheat rotation. *Journal of Research Punjab Agricultural University*, 22:243-52.
- Singh M, Reddy SR, Singh VP, Rupa TR (2007). Phosphorus availability to rice (*Oriza sativa* L.) wheat (*Triticum estivum* L.) in a Vertisol after eight years of inorganic and organic fertilizer additions. *Bioresource Technology*, 98: 1474-1481.
- Singh NP, Sachan RS, Pandey PC, Bisht PS (1999). Effect of a decade long-term fertilizer and manure application on fertility and productivity of rice-wheat system in a mollisols. *Journal of Indian Society of Soil Science*, 47(1): 72-80.
- Sinha SK, Singh VN, Singh KP (1997). Effect of Continuous Application of Manures and Fertilisers on Available Nutrients in an Alluvial Soil. *Journal of Research-birsa Agricultural University*, 9:163-6.
- Subbiah BV, Asija GL (1956). A Rapid procedure for estimation of available nitrogen in rice soils. *Current Science*, 25: 259-260.
- Subehia SK, Sepehya S (2012). Effect of Integrated Nutrient Management on rice-wheat yield and soil properties in a North Western Himalayan Region. *Indian Journal of soil Conservation*, 40(2): 135-140.
- Tillman AK (1998). The influence of long-term organic and mineral fertilization on soil organic matter. *Soil Water Resource*, 3(2): 41-51.
- Tyagi VV (1989). Soil fertility status and physic chemical status of an aquic Hapludoll as influenced by 16 years of continuous cropping and fertilization. Thesis M.Sc. (Ag.) submitted to G. B. P. U. A & T Pantnagar. (U.S. Nagar), 10-24.
- Urkurkar JS, Tewari S, Bajpai RK (2010). Influence of long term use of organic and inorganic manures on soil fertility and sustainable productivity of rice (*Oryza sativa*) and wheat (*Triticum aestivum*) in Inceptisols. *Indian Journal of Agricultural Science*, 80(3): 208-212.
- Walkley A, Black IA (1934). An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*, 37(1): 29-38.
- Yadav DS, Kumar A (2009). Long-term effect of nutrient management on soil health and productivity of rice (*Oryza Sativa*)-wheat (*Triticum Aestivum*) system. *Indian Journal of Agronomy*, 54(1): 15-23.
- Yadhuvanshi NPS (2003). Substantiation of inorganic fertilizer by organic manure and the effect on soil fertility in a rice-wheat rotation on reclaimed sodic soil in India. *Indian Journal of Agricultural Science*, 140: 161-168.